



Note on the Carleman's inequality and Hardy's inequality[☆]

Lü Zhongxue^{a,*}, Gao Youcai^b, Wei Yuxiang^a

^a School of Mathematical Science, Xuzhou Normal University, Xuzhou, 221116, PR China

^b Daqing Normal University, Daqing, 163712, PR China

ARTICLE INFO

Article history:

Received 2 March 2009

Received in revised form 3 September 2009

Accepted 4 September 2009

Keywords:

Hardy's inequality

Hardy–Littlewood's inequality

Carleman's inequality

ABSTRACT

In this article, using the properties of power mean and induction, new strengthened Carleman's inequality and Hardy's inequality are obtained. We also give an answer to the conjectures proposed by X. Yang in the literature Yang (2001) [5].

© 2009 Elsevier Ltd. All rights reserved.

1. Introduction

The following Carleman's inequality is well known (see [1])

$$\sum_{n=1}^{\infty} (a_1 a_2 \cdots a_n)^{1/n} < e \sum_{n=1}^{\infty} a_n \quad (1)$$

where $a_n \geq 0$, $n = 1, 2, \dots$, and $0 < \sum_{n=1}^{\infty} a_n < \infty$.

By Hardy (see, [1, Theorem 349]), the Carleman's inequality was generalized as follows:

$$\sum_{n=1}^{\infty} \lambda_{n+1} (a_1^{\lambda_1} a_2^{\lambda_2} \cdots a_n^{\lambda_n})^{1/\Lambda_n} < e \sum_{n=1}^{\infty} \lambda_n a_n \quad (2)$$

where $a_n \geq 0$, $\lambda_n > 0$, $\Lambda_n = \sum_{m=1}^n \lambda_m$, $n = 1, 2, \dots$, and $0 < \sum_{n=1}^{\infty} \lambda_n a_n < \infty$.

In recent years, Yang [2], Yan and Sun [3], Xie and Zhong [4], Yang [5], and Dragomir and Ho Kim [6] gave some distinct improvements of (1) or (2), respectively.

Yang [5] proposed the following conjectures:

Conjecture 1. Let $x \geq 0$. Then the following equality holds

$$\left(1 + \frac{1}{x}\right)^x = e \left[1 - \sum_{k=1}^{\infty} \frac{b_k}{(1+x)^k}\right],$$

where $b_1 = \frac{1}{2}$, $b_2 = \frac{1}{24}$, $b_3 = \frac{1}{48}$, $b_4 = \frac{73}{5760}$, $b_5 = \frac{11}{1280}$ and $b_6 = \frac{1945}{580608}$.

Conjecture 2. Is $b_k > 0$, for $k \geq 7$?

In this note, we give an answer on these conjectures. New, strengthened Carleman's and Hardy's inequalities are also obtained.

[☆] This work is supported the Natural Science Foundation of China (No.10771181) and NSF of Jiangsu Higher Education Bureau (No. 07KJD110206).

* Corresponding author.

E-mail address: lvzx1@tom.com (L. Zhongxue).

2. Some lemmas

We will use the following lemmas.

Lemma 2.1 ([6]). Let $x > 0$, then

$$\left(1 + \frac{1}{x}\right)^x = e \left[1 - \sum_{k=1}^{\infty} \frac{b_k}{(1+x)^k}\right], \quad (3)$$

where $b_1 = \frac{1}{2}$, $b_{n+1} = \frac{1}{n+1} \left(\frac{1}{n+2} - \sum_{k=1}^n \frac{b_k}{n+2-k}\right)$, $n = 1, 2, \dots$

Lemma 2.2. Let $x > 0$, $m \in \mathbb{N}$, then

$$\left(1 + \frac{1}{x}\right)^x < e \left[1 - \sum_{k=1}^m \frac{b_k}{(1+x)^k}\right], \quad (4)$$

where $b_1 = \frac{1}{2}$, $b_{n+1} = \frac{1}{n+1} \left(\frac{1}{n+2} - \sum_{k=1}^n \frac{b_k}{n+2-k}\right)$, $n = 1, 2, m-1$.

Proof. By Lemma 2.1, we only prove that $b_n > 0$, $n \in \mathbb{N}$. By induction, because $b_1 = \frac{1}{2} > 0$, we will prove that: if $b_t > 0$, as $n = t-1$, $t \in \mathbb{N}$ and $t \geq 2$, then $b_{t+1} > 0$, as $n = t$.

In fact, when $n = t$ ($t \in \mathbb{N}$ and $t \geq 2$),

$$\begin{aligned} b_{t+1} &= \frac{1}{t+1} \left(\frac{1}{t+2} - \sum_{k=1}^t \frac{b_k}{t+2-k} \right) \\ &= \frac{1}{t+1} \left(\frac{1}{t+2} - \sum_{k=1}^{t-1} \frac{b_k}{t+2-k} - \frac{b_t}{2} \right) \\ &= \frac{1}{t+1} \left[\left(\frac{1}{t+2} - \frac{1}{2t(t+1)} \right) - \sum_{k=1}^{t-1} \left(\frac{1}{t+2-k} - \frac{1}{2t(t+1-k)} \right) b_k \right] \\ &= \frac{1}{t+1} \left[\left(\frac{t+1}{t+2} - \frac{1}{2t} \right) \frac{1}{t+1} - \sum_{k=1}^{t-1} \left(\frac{t+1-k}{t+2-k} - \frac{1}{2t} \right) \frac{b_k}{t+1-k} \right] \\ &> \frac{1}{t+1} \left[\left(\frac{t+1}{t+2} - \frac{1}{2t} \right) \frac{1}{t+1} - \left(\frac{t+1}{t+2} - \frac{1}{2t} \right) \sum_{k=1}^{t-1} \frac{b_k}{t+1-k} \right] \\ &= \frac{t}{t+1} \left(\frac{t+1}{t+2} - \frac{1}{2t} \right) \left[\frac{1}{t} \left(\frac{1}{t+1} - \sum_{k=1}^{t-1} \frac{b_k}{t+1-k} \right) \right] \\ &= \frac{t}{t+1} \left(\frac{t+1}{t+2} - \frac{1}{2t} \right) b_t. \end{aligned}$$

By induction, we have $b_n > 0$, $n \in \mathbb{N}$. The lemma is proved. \square

Remark 2.3. Lemmas 1.1 and 2.2 have given an answer of Conjectures 1 and 2.

By Lemma 2.2, we have the following corollary.

Corollary 2.4. Let $a_n \geq 0$, $m \in \mathbb{N}$, $0 < \lambda_{n+1} \leq \lambda_n$, $A_n = \sum_{m=1}^n \lambda_n$ satisfying $A_n \geq 1$, then

$$\left(1 + \frac{\lambda_n}{A_n}\right)^{A/\lambda} < e \left[1 - \sum_{k=1}^m \frac{\lambda_k b_k}{(\lambda_n + A_n)^k}\right], \quad (5)$$

where $b_1 = \frac{1}{2}$, $b_{n+1} = \frac{1}{n+1} \left(\frac{1}{n+2} - \sum_{k=1}^n \frac{b_k}{n+2-k}\right)$, $n = 1, 2, m-1$.

3. Main results

Now we introduce the main results.

Theorem 3.1. Let $0 < \lambda_{n+1} \leq \lambda_n$, $\Lambda_n = \sum_{m=1}^n \lambda_n$ satisfying $\Lambda_n \geq 1$, $a_n \geq 0$ ($n \in \mathbb{N}$), $0 < p \leq 1$, $m \in \mathbb{N}$ and $0 < \sum_{n=1}^m \lambda_n a_n < \infty$. Then

$$\sum_{n=1}^{\infty} \lambda_{n+1} (a_1^{\lambda_1} a_2^{\lambda_2} \cdots a_n^{\lambda_n})^{1/\Lambda_n} < \frac{e^p}{p} \sum_{n=1}^{\infty} \left(1 - \sum_{k=1}^m \frac{\lambda_n^k b_k}{(\lambda_n + \Lambda_n)^k} \right)^p \lambda_n a_n^p \Lambda_n^{p-1} \left(\sum_{s=1}^n \lambda_s (c_s a_s)^p \right)^{(1-p)/p} \quad (6)$$

where $c_s^{\lambda_n} = \Lambda_{n+1}^{\lambda_n} / \Lambda_n^{\lambda_{n-1}}$, $b_1 = \frac{1}{2}$, $b_{n+1} = \frac{1}{n+1} \left(\frac{1}{n+2} - \sum_{k=1}^n \frac{b_k}{n+2-k} \right)$, $n = 1, 2, m-1$.

Proof. By the power mean inequality, we have

$$\begin{aligned} \sum_{n=1}^{\infty} \lambda_{n+1} (a_1^{\lambda_1} a_2^{\lambda_2} \cdots a_n^{\lambda_n})^{1/\Lambda_n} &= \sum_{n=1}^{\infty} \lambda_{n+1} \frac{(c_1 a_1)^{\lambda_1/\Lambda_n} (c_2 a_2)^{\lambda_2/\Lambda_n} \cdots ((c_n a_n)^{\lambda_n/\Lambda_n})^{1/\Lambda_n}}{(c_1^{\lambda_1} c_2^{\lambda_2} \cdots c_n^{\lambda_n})^{1/\Lambda_n}} \\ &\leq \sum_{n=1}^{\infty} \frac{\lambda_{n+1}}{(c_1^{\lambda_1} c_2^{\lambda_2} \cdots c_n^{\lambda_n})^{1/\Lambda_n}} \left(\frac{1}{\Lambda_n} \sum_{m=1}^n \lambda (c_m a_m)^p \right)^{1/p} \end{aligned} \quad (7)$$

For $c_m > 0$, $m = 1, 2, \dots, n$.

By using the Hardy–Littlewood inequality (see [7,8]), we have

$$\begin{aligned} \left(\frac{1}{\Lambda_n} \sum_{m=1}^n \lambda_m (c_m a_m)^p \right)^{1/p} &\leq \frac{1}{\Lambda_n} \left(\sum_{m=1}^n \lambda_m (c_m a_m)^p \right)^{1/p} \\ &\leq \frac{1}{p \Lambda_n} \sum_{m=1}^n \lambda_m (c_m a_m)^p \left(\sum_{s=1}^m \lambda_s (c_s a_s)^p \right)^{(1-p)/p}. \end{aligned} \quad (8)$$

For $\Lambda_n \geq 1$, $0 < p \leq 1$.

Choosing $c_1^{\lambda_1} c_2^{\lambda_2} \cdots c_n^{\lambda_n} = \Lambda_{n+1}^{\Lambda_n}$ ($n \in \mathbb{N}$), and setting $\Lambda_0 = 0$, from $\lambda_{n+1} \leq \lambda_n$, it follows that

$$c_n = \left[\frac{\Lambda_{n+1}^{\Lambda_n}}{\Lambda_n^{\Lambda_{n-1}}} \right]^{1/\lambda_n} = \left[1 + \frac{\lambda_{n+1}}{\Lambda_n} \right]^{\Lambda_n/\lambda_n} \cdot \Lambda_n \leq \left[1 + \frac{\lambda_n}{\Lambda_n} \right]^{\Lambda_n/\lambda_n} \cdot \Lambda_n. \quad (9)$$

From (7)–(9), it follows that

$$\sum_{n=1}^{\infty} \lambda_{n+1} (a_1^{\lambda_1} a_2^{\lambda_2} \cdots a_n^{\lambda_n})^{1/\Lambda_n} \leq \frac{1}{p} \sum_{m=1}^{\infty} \left[1 + \frac{\lambda_m}{\Lambda_m} \right]^{p \Lambda_m / \lambda_m} \cdot \lambda_m \Lambda_m^{p-1} a_m^p \left(\sum_{s=1}^m \lambda_s (c_s a_s)^p \right)^{(1-p)/p}.$$

Hence by the above inequality and Lemma 2.2, we have

$$\sum_{n=1}^{\infty} \lambda_{n+1} (a_1^{\lambda_1} a_2^{\lambda_2} \cdots a_n^{\lambda_n})^{1/\Lambda_n} < \frac{e^p}{p} \sum_{n=1}^{\infty} \left(1 - \sum_{k=1}^m \frac{\lambda_n^k b_k}{(\lambda_n + \Lambda_n)^k} \right)^p \lambda_n a_n^p \Lambda_n^{p-1} \left(\sum_{s=1}^n \lambda_s (c_s a_s)^p \right)^{(1-p)/p}.$$

Thus Theorem 3.1 is proved. \square

Setting $p = 1$ in Theorem 3.1, then we get an extension of the strengthened Hardy's inequality as following.

Corollary 3.2. Let $0 < \lambda_{n+1} \leq \lambda_n$, $\Lambda_n = \sum_{m=1}^n \lambda_n$ satisfying $\Lambda_n \geq 1$, $a_n \geq 0$ ($n \in \mathbb{N}$), $m \in \mathbb{N}$ and $0 < \sum_{n=1}^m \lambda_n a_n < \infty$. Then

$$\sum_{n=1}^{\infty} \lambda_{n+1} (a_1^{\lambda_1} a_2^{\lambda_2} \cdots a_n^{\lambda_n})^{1/\Lambda_n} < e \sum_{n=1}^{\infty} \left(1 - \sum_{k=1}^m \frac{\lambda_n^k b_k}{(\lambda_n + \Lambda_n)^k} \right) \lambda_n a_n \quad (10)$$

where $c_s^{\lambda_n} = \Lambda_{n+1}^{\lambda_n} / \Lambda_n^{\lambda_{n-1}}$, $b_1 = \frac{1}{2}$, $b_{n+1} = \frac{1}{n+1} \left(\frac{1}{n+2} - \sum_{k=1}^n \frac{b_k}{n+2-k} \right)$, $n = 1, 2, m-1$.

Setting $\lambda_n = 1$ in Theorem 3.1, then we also get an extension of the strengthened Carleman's inequality as the following.

Corollary 3.3. Let $a_n \geq 0$ ($n \in \mathbb{N}$), $0 < p \leq 1$, $m \in \mathbb{N}$ and $0 < \sum_{n=1}^m \lambda_n a_n < \infty$. Then

$$\sum_{n=1}^{\infty} (a_1 a_2 \cdots a_n)^{1/n} < \frac{e^p}{p} \sum_{n=1}^{\infty} \left(1 - \sum_{k=1}^m \frac{\lambda_n^k b_k}{(1+n)^k} \right)^p a_n^p n^{p-1} \left(\sum_{s=1}^n (c_s a_s)^p \right)^{(1-p)/p} \quad (11)$$

where $c_s = (1 + 1/s)^s \cdot s$, $b_1 = \frac{1}{2}$, $b_{n+1} = \frac{1}{n+1} \left(\frac{1}{n+2} - \sum_{k=1}^n \frac{b_k}{n+2-k} \right)$, $n = 1, 2, m-1$.

Similarly to the proof of [Theorem 3.1](#), we can prove the generalized version of Hardy's inequality as the following theorem:

Theorem 3.4. Let $0 < \lambda_{n+1} \leq \lambda_n$, $\Lambda_n = \sum_{m=1}^n \lambda_m$ satisfying $\Lambda_n \geq 1$, $a_n \geq 0$ ($n \in \mathbb{N}$), $m \in \mathbb{N}$ and $0 < \sum_{n=1}^m \lambda_n a_n^t < \infty$ for $0 < p \leq t < \infty$, Then

$$\sum_{n=1}^{\infty} \lambda_{n+1} (a_1^{\lambda_1} a_2^{\lambda_2} \cdots a_n^{\lambda_n})^{t/\Lambda_n} < \frac{te^{p/t}}{p} \sum_{n=1}^{\infty} \left(1 - \sum_{k=1}^m \frac{\lambda_n^k b_k}{(\lambda_n + \Lambda_n)^k} \right)^{p/t} \lambda_n a_n^p \Lambda_n^{(p-t)/t} \left(\sum_{s=1}^n \lambda_s (c_s a_s)^p \right)^{(t-p)/p} \quad (12)$$

where $c_s^{\lambda_n} = \Lambda_n^{\lambda_n} / \Lambda_n^{\lambda_{n-1}}$, $b_1 = \frac{1}{2}$, $b_{n+1} = \frac{1}{n+1} (\frac{1}{n+2} - \sum_{k=1}^n \frac{b_k}{n+2-k})$, $n = 1, 2, m-1$.

The proof is almost the same as in proving [Theorem 3.1](#). We here only need to note that

$$(\alpha_1^{q_1} \alpha_2^{q_2} \cdots \alpha_n^{q_n})^t \leq \left(\sum_{m=1}^n q_m a_m^p \right)^{t/p}, \quad p, t \geq 0$$

where $\alpha_m \geq 0$, $0 < p \leq t < \infty$, $q_m > 0$, and $\sum_{m=1}^n q_m = 1$ ($m \in \mathbb{N}$).

Setting $p = 1$ in [Theorem 3.4](#), then we get an extension of the strengthened Hardy's inequality as the following.

Corollary 3.5. Let $0 < \lambda_{n+1} \leq \lambda_n$, $\Lambda_n = \sum_{m=1}^n \lambda_m$ satisfying $\Lambda_n \geq 1$, $a_n \geq 0$ ($n \in \mathbb{N}$), $m \in \mathbb{N}$ and $0 < \sum_{n=1}^m \lambda_n a_n^t < \infty$ for $1 \leq t < \infty$. Then

$$\sum_{n=1}^{\infty} \lambda_{n+1} (a_1^{\lambda_1} a_2^{\lambda_2} \cdots a_n^{\lambda_n})^{t/\Lambda_n} < te^{1/t} \sum_{n=1}^{\infty} \left(1 - \sum_{k=1}^m \frac{\lambda_n^k b_k}{(\lambda_n + \Lambda_n)^k} \right)^{1/t} \lambda_n a_n \Lambda_n^{(1-t)/t} \left(\sum_{s=1}^n \lambda_s (c_s a_s) \right)^{(t-1)}. \quad (13)$$

where $c_s^{\lambda_n} = \Lambda_n^{\lambda_n} / \Lambda_n^{\lambda_{n-1}}$, $b_1 = \frac{1}{2}$, $b_{n+1} = \frac{1}{n+1} (\frac{1}{n+2} - \sum_{k=1}^n \frac{b_k}{n+2-k})$, $n = 1, 2, m-1$.

Setting $\lambda_n = 1$ in [Theorem 3.4](#), then we also get an extension of the strengthened Carleman's inequality as the following.

Corollary 3.6. Let $a_n \geq 0$ ($n \in \mathbb{N}$), $m \in \mathbb{N}$ and $0 < \sum_{n=1}^m a_n^t < \infty$ for $0 < p \leq t < \infty$, Then

$$\sum_{n=1}^{\infty} (a_1 a_2 \cdots a_n)^{t/n} < \frac{te^{p/t}}{p} \sum_{n=1}^{\infty} \left(1 - \sum_{k=1}^m \frac{b_k}{(1+n)^k} \right)^{p/t} a_n^p n^{(p-t)/t} \left(\sum_{s=1}^n (c_s a_s)^p \right)^{(t-p)/p} \quad (14)$$

where $c_s^t = (1 + 1/s)^s \cdot s$, $b_1 = \frac{1}{2}$, $b_{n+1} = \frac{1}{n+1} (\frac{1}{n+2} - \sum_{k=1}^n \frac{b_k}{n+2-k})$, $n = 1, 2, m-1$.

References

- [1] G.H. Hardy, J.E. Littlewood, G. Polya, Inequalities, Cambridge Univ. Press, London, 1952.
- [2] Yang Bicheng, Note on Hardy's inequality, J. Math. Anal. Appl. 234 (1999) 717–722.
- [3] P. Yan, G. Sun, A strengthened Carleman's inequality, J. Math. Anal. Appl. 240 (1999) 290–293.
- [4] Z. Xie, Y. Zhong, A best approximation for constant e and an improvement to Hardy's inequality, J. Math. Anal. Appl. 234 (1999) 717–722.
- [5] X. Yang, A Carleman's inequality, J. Math. Anal. Appl. 253 (2001) 691–694.
- [6] S. Dragomir, Ho Kim, The strengthened Hardy's inequality and their new generalizations, Fac. Sci. Math. 20 (2) (2006) 39–49.
- [7] G.S. Davis, G.M. Peterson, On an inequality of Hardy's (II), Quart. J. Math. (Oxford) 15 (1964) 35–40.
- [8] J. Németh, Generalizations of the Hardy–Littlewood inequality, Acta. Sci. Math. (Szeged) 32 (1971) 295–299.